



ASO VISUAL ABSTRACT

ASO Visual Abstract: Glissonian Approach During Robotic Mesohepatectomy for Recurrent Colorectal Liver Metastasis

Marcel Aufran C. Machado, MD, FACS, Bruno V. Mattos, MD, Murillo Macedo Lobo Filho, MD, and Fabio Makdissi, MD


Nove de Julho Hospital, São Paulo, Brazil

Hepatectomy is the standard treatment for colorectal liver metastases (CLM); however, the high recurrence rate of CLM is a problem and may be related to missing metastases. Repeat hepatectomy may offer favorable survival but is technically demanding, therefore minimally invasive repeat hepatectomy has been used in only a few patients. The purpose of this manuscript is to describe the importance and feasibility of the Glissonian approach during robotic redo hepatectomy. We present a video of a

robotic mesohepatectomy for recurrent CLM (<https://doi.org/10.1245/s10434-022-12331-4>). The robotic approach is safe and feasible for mesohepatectomy and may be a good indication for re-hepatectomy after open surgery. The intrahepatic Glissonian approach is useful for anatomical central resection. We conclude that the Glissonian approach is useful even during robotic liver resections, especially redo hepatectomies. This video may help oncological surgeons to perform this complex procedure.

Glissonian Approach During Robotic Mesohepatectomy for Recurrent Colorectal Liver Metastasis

Repeat Liver Resection after Multiple Open Procedures



Glissonian Approach

Purpose: Anatomical Resection after Previous Liver Resection

Glissonian Approach

Anatomical Resection **Liver-Sparing technique**

Conclusion: Use of Glissonian Approach is Useful Even During Robotic Liver Resections

Machado et al. *Ann Surg Oncol*.
Visual Abstract @AnnSurgOncol

ANNALS OF SURGICAL ONCOLOGY

DISCLOSURE Marcel Aufran Machado, Bruno V. Mattos, Murillo Macedo Lobo Filho, and Fabio Makdissi declare no conflicts of interest.