ASO VISUAL ABSTRACT


ASO Visual Abstract: Robotic Isolated Pancreaticojejunostomy After Pancreatoduodenectomy: A Good Choice for Young Patients with Expected Long-Term Survival

Marcel Autran C. Machado, MD, FACS, Bruno V. Mattos, MD, Murillo Macedo Lobo Filho, MD, Augusto C. Carvalho, MD, Fabio Makdissi, MD, and Marcel C. C. Machado, MD, FACS

Nove de Julho Hospital, São Paulo, Brazil

Isolated pancreatojejunostomy after pancreatoduodenectomy (PD) can reduce the severity of pancreatic fistula and may be a good option for young patients with benign or low-grade neoplasms as well as for patients with a soft pancreas and high risk of clinically relevant postoperative pancreatic fistula (POPF). It may increase the surgical reconstruction

time, but not the overall time, because the resection phase usually is easier in those cases. This procedure can be performed safely with the robotic platform. This video can help oncologic surgeons perform this complex procedure (https://doi.org/10.1245/s10434-025-17037-x).



Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© Society of Surgical Oncology 2025

M. A. C. Machado, MD, FACS e-mail: dr@drmarcel.com.br

Published online: 22 March 2025